



A priori bounds for the nonlinear Parabolic Anderson Model

Hendrik Weber University of Munster

Hendrik Weber is a W3 professor at University of Munster since September 2022. Previously, he was a professor of probability at University of Bath and a Reader at the University of Warwick. He is a recipient of a 2022 ERC Consolidator Grant, the 2017 Philip Leverhulme Prize, and the 2016 Rollo Davidson Prize.



Abstract: We show a priori bounds that exclude finite time explosion for the solution of the nonlinear (or generalised) Parabolic Anderson model (gPAM) $(\partial_t - \Delta)u = \sigma(u)\xi$ in the framework of Hairer's regularity structures. gPAM is the archetype of an equation that can be treated using regularity structures and it was one of the first equations for which a local solution theory was developed using this theory. While global existence in the linear case $\sigma(u) = u$ follows from Hairer's original work [Hairer, Inventiones 2014] and also [Gubinelli, Imkeller, Perkowski FoM, Pi 2015], the case of non-linear σ is significantly more challenging and has remained open so far. Our results apply to noise terms ξ of (ir)regularity $> -1 - \kappa$ for some $\kappa > 0$, including in particular the case of 2-dimensional spacial white noise. As a corollary we obtain global existence and the existence of an invariant measure for the dynamic Sine-Gordon model on the two-dimensional torus in the regime $\beta^2 \in (4\pi, (1 + \kappa)4\pi)$.

This is joint work with Guilherme Feltes ((Münster) and Ajay Chandra (Imperial College).

讲座时间:

2023. 12. 18周一下午17:00-18:00

会议地点: ZOOM会议室 会议ID: 354 143 7366 密码: 123456

主办单位:

中科院数学与系统科学研究院应用数学所

北京理工大学数学与统计学院